

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 45/50, B01J 31/18, C07C 47/277, 47/23, 45/49	A1	(11) International Publication Number: WO 95/06627 (43) International Publication Date: 9 March 1995 (09.03.95)
(21) International Application Number: PCT/US94/09461 (22) International Filing Date: 26 August 1994 (26.08.94) (30) Priority Data: 08/116,235 2 September 1993 (02.09.93) US (71) Applicant: E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: AYERS, Timothy, Allen; 416 Brandywine Boulevard, Wilmington, DE 19803 (US). RAJANBABU, Thaliyil, V.; 2304 Ramblewood Drive, Wilmington, DE 19810 (US). (74) Agents: GEIGER, Kathleen, W. et al.; E.I. du Pont de Nemours and Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ENANTIOSELECTIVE HYDROFORMYLATION (57) Abstract Enantioselective hydroformylation of vinyl compounds using a catalyst composition of a chiral carbohydrate phosphorous ligand with a Rh, Pt, Co or Ir metal, to produce chiral 2-substituted propanals, wherein the phosphorous of the ligand is substituted with electron withdrawing groups.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MD	Republic of Moldova	TJ	Tajikistan
DE	Germany	MG	Madagascar	TT	Trinidad and Tobago
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain	MN	Mongolia	US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE

ENANTIOSELECTIVE HYDROFORMYLATION

FIELD OF THE INVENTION

The present invention relates to enantioselective
5 hydroformylation of vinyl compounds $R-CH=CH_2$ to produce
chiral nonracemic 2-substituted 1-propanals of the
formula $R-CH(CHO)CH_3$; wherein a catalyst composition of
the reaction comprises Rh, Pt, Co or Ir, and a chiral,
nonracemic carbohydrate phosphorous ligand wherein the
10 phosphorus is substituted with electron withdrawing
groups.

BACKGROUND OF THE INVENTION

It is known in the art that optically active
diphosphetes derived from carbohydrates can be used as
15 ligands for the metals Rh and Pt in the asymmetric
hydroformylation of vinyl arenes. The major problem
that has plagued the well-known Pt-catalyzed
hydroformylations, however, is the unfavorable branched
to linear ratios which reduce overall selectivity. The
20 Rh-catalyzed reactions generally give good branched to
linear ratios but enantioselectivity is poor. A series
of metal/ligand combinations which overcome these
limitations is the subject of this application.

Ojima, I. and Hirai, K., "Asymmetric Hydro-
25 silylation and Hydrocarbonylation" in Asymmetric
Synthesis; Morrison, J. D., Ed.; Academic Press,
Orlando, Florida, 1985; pp 103-146 and Jackson, W. R.
and Lovel, C. G., Aust. J. Chem., 35, 2069-75 (1982)
describe low asymmetric induction in the hydro-
30 formylation of vinyl acetate in the presence of a
phosphinite derived from tartaric acid.

German patents DD280,473, DD275,623 and DD275,671
and references Selke et al., J. Mol. Cat., 37, 213-225
(1986) and Selke, J. Organometal. Chem., 370, 249-256
35 (1989) disclose related rhodium carbohydrate catalysts


mainly for enantioselective hydrogenation reactions, but no teaching is provided which demonstrates or enables effective enantioselective hydroformylation, and the importance of electronic effects of these ligands is not suggested.

Ligands similar to those used in the present invention are disclosed in RajanBabu, T. V. and Casalnuovo, A. L., *J. Am. Chem. Soc.*, 114, 6265-6266 (1992) and U.S. Patent 5,175,335, for use in other catalyst compositions for asymmetric hydrocyanation.

High enantioselectivities are most often explained on the basis of steric arguments, see for example, Brown, J. M., *Chem. Soc. Rev.*, 25 (1993), and references cited therein. For a recent example of the application of classical and widely used steric approach to design of enantioselective catalysts see: Trost, B. M. et al. *J. Am. Chem. Soc.* 114, 9327 (1992) and references cited therein. Recently, electronic effects are described as being effective in enhancing enantioselectivity in manganese-catalyzed oxidation, Jacobsen, E. N. et al., *J. Am. Chem. Soc.* 113, 6703, (1991), and in nickel-catalyzed hydrocyanations, RajanBabu, T. V. and Casalnuovo, A. L., *J. Am. Chem. Soc.*, 114, 6265-6266 (1992). No teaching of electronic effects in the enantioselective hydroformylation reaction have been described.

SUMMARY OF THE INVENTION

The present invention provides a process for enantioselective hydroformylation comprising: reacting a vinyl compound of formula I



I

with a source of CO and H₂, in the presence of a catalyst composition comprising one or more metals


3

selected from the group consisting of Co, Rh, Ir, and Pt, and a chiral, nonracemic ligand of formula II



II

to produce a nonracemic hydroformylated product of formula III



III

5 wherein:

R is a C₁ to C₂₀ carboalkoxy, a C₁ to C₄₀ hydrocarbyl, or a C₁ to C₄₀ heterocyclic radical; each optionally substituted with one or more halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups;

10

each R¹ is an electron-withdrawing group comprising an aromatic hydrocarbyl substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkylsulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups; or a heteroaromatic optionally substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups;

15

20

R² is a C₄ to C₄₀ dideoxy carbohydrate optionally substituted with one or more

hydrocarbyl, halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups.

The present invention also provides a novel catalyst composition comprising one or more metals selected from the group consisting of Co, Rh, Ir and Pt, and one or more chiral nonracemic ligands of formula II



II

wherein:

each R^1 is an electron-withdrawing group comprising an aromatic hydrocarbyl or heteroaromatic, optionally substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups; and


R^2 is a C_4 to C_{40} dideoxy carbohydrate; optionally substituted with one or more hydrocarbyl, halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups.

DETAILED DESCRIPTION OF THE INVENTION

The process of the instant invention, whereby enantioselective hydroformylation is accomplished by reacting a vinyl compound of the formula $R-CH=CH_2$ with hydrogen and carbon monoxide in the presence of a chiral, nonracemic, metal (Rh, Pt, Ir or Co) hydroformylation catalyst, is useful, for example, to produce 2-substituted-1-propanals. For example, vinyl arenes are precursors to nonsteroidal, anti-inflammatory drugs such as ibuprofen and naproxen. The novel catalyst compositions of the instant invention, comprising a metal and chiral, nonracemic diphosphinite ligand derived from a carbohydrate diol, are useful for

accomplishing the above-described enantioselective hydroformylation reactions.

The enantioselective hydroformylation reaction of the invention is performed by reacting a vinyl compound of the formula $R-CH=CH_2$ with hydrogen and carbon monoxide at pressures of about 100-2400 psi (1 psi = 6.9 kPa) in the presence of a chiral, nonracemic, metal (Rh, Pt, Ir or Co) hydroformylation catalyst. These reactions produce chiral, nonracemic, 2-substituted-1-propanals of the formula $R-CH(CHO)CH_3$, herein called branched aldehydes. Undesired products, 3-substituted-1-propanals are also generated, herein called linear aldehydes. The selectivity of this reaction will be reported herein by the ratio of branched aldehyde (b) to the linear aldehyde (l).




By the term "carbohydrate", Applicants mean the class of organic compounds comprising the general formula $(CH_2O)_n$, wherein n is equal to or greater than four. The carbohydrate-derived ligands of the invention are derived from C_4 to C_{40} carbohydrates including monosaccharides, disaccharides and oligosaccharides.

By the term "heterocycle", Applicants mean a cyclic carbon compound containing at least one oxygen, nitrogen or sulfur atom in the ring.


By the term "hydrocarbyl", Applicants include all alkyl, aryl, aralkyl or alkylaryl carbon substituents, either straight-chained, cyclic, or branched, accordingly substituted with hydrogen. The term "hydrocarbyl" as used herein includes both aromatic and nonaromatic hydrocarbyls.

By the term electron-withdrawing group, Applicants include those groups that have σ -values (any σ -values such as σ_p , σ_m , σ_i , σ_R , etc.) greater than or equal to 0.1 (as defined by the Hammett equation, see for example March, J. Advanced Organic Chemistry, Reactions, Mechanisms, and Structure, 4th ed.; 1992, Wiley: New York, pp 278-286).


In describing a carbohydrate group of the formula $O-R^2-O$, as it appears within the ligand of the present disclosure, the group R^2 is named by using the prefix "dideoxy" with the name of the parent diol of the formula $HO-R^2-OH$. For example, the name 2,3-dideoxy-glucose refers to the group:



and accordingly, the corresponding carbohydrate group $O-R^2-O$ is:



The suffix -ose- when used in combination with carbohydrate root names, shall include those compounds wherein the OH groups are protected as ethers or esters. By this definition, for example, the glucopyranoside structure shown below is termed "a glucose"



wherein Ac is an acetyl.

By the term "chiral", Applicants mean "existing as a pair of enantiomers". These enantiomers, where the chiral centers are designated the R and S isomers, are nonsuperimposable mirror images of one another. A chiral material may either contain an equal amount of the R and S isomers in which case it is called "racemic" or it may contain inequivalent amounts of R and S isomer in which case it is called "optically active", or "nonracemic".

By the term "enantiomeric excess" ("ee"), Applicants mean the absolute difference between the percent of R enantiomer and the percent of S enantiomer of an optically active compound. For example, a compound which contains 75% S isomer and 25% R isomer will have an enantiomeric excess of 50%.

By the term "enantioselective" Applicants mean the ability to produce a product in an optically active form.

The substrates of the invention are described by the formula $R-CH=CH_2$, where R may be any C_1 to C_{20} carboalkoxy, C_1 to C_{40} hydrocarbyl or C_1 to C_{40} heterocyclic radical; each of which may be substituted with one or more halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups. Examples of R include, but are not limited to, phenyl, substituted phenyl, polyaromatic (e.g., naphthyl, anthryl), substituted polyaromatic, acetoxy, alkyl and substituted alkyl. Representative examples of substrates used in the invention include, but are not limited to, 2-vinylnaphthalene, 6-methoxy-2-

vinyl naphthalene, vinyl acetate, 4-isobutylstyrene, 4-methylstyrene and styrene.

The vinyl substrates of the invention may be made by methods which are well-known in the art e.g.,

- 5 Organometallics, 10, 1183-1189 (1991), which is hereby incorporated by reference. Many substrates are also available commercially.

The chiral, nonracemic carbohydrate diphosphinite ligands of the invention are defined as

- 10 $(R^1)_2-P-O-R^2-O-P-(R^1)_2$, wherein R^1 may be an aromatic hydrocarbyl substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups; or a heteroaromatic optionally substituted with
15 one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups. These ligands are unique in this art in that the R^1 group of the ligand is an electron-withdrawing group as defined herein.
20 Applicants have discovered that electron-withdrawing groups generally lead to high selectivity in the hydroformylation method of this invention.





- For all embodiments of the Applicants' invention, the chiral, nonracemic, metal hydroformylation catalyst
25 composition comprises a chiral, nonracemic, carbohydrate diphosphinite ligand and a source of one or more of the metals Rh, Pt, Co and Ir. Suitable sources of the rhodium, platinum, cobalt, and iridium include, but are not limited to, the metal halides, olefin complexes,
30 acetoacetates, and carbonyls. Metal compounds that contain ligands which can be displaced by the chiral carbohydrate phosphorus ligand are a preferred source of the metal. In the case, for example, of rhodium (I) intermediates, $(COD)_2RhX$ species (COD is 1,5-cycloocta-
35 diene) are the precursors of choice, with the counterion

X being tetrafluoroborate (BF_4), antimony hexafluoride (SbF_6), or trifluoromethanesulfonate (OTf); although other counterions such as tetraphenylborate (BPh_4) and perchlorate (ClO_4) would also be suitable. Chiral
5 iridium compounds can be prepared from $[(\text{COD})\text{IrCl}]_2$. Platinum compounds can be synthesized from dichlorobis(benzo-nitrile)platinum (II). Cobalt compounds can be prepared from cobalt carbonyl. Rhodium is the preferred metal.

10 The catalyst composition also employs a ligand comprising a chiral, nonracemic diphosphinite of the formula $(\text{R}^1)_2\text{-P-O-R}^2\text{-O-P-(R}^1)_2$, wherein the R^2 is a C_4 to C_{40} dideoxycarbohydrate, optionally substituted with one or more hydrocarbyl, halo, alkoxy, carboalkoxy, hydroxy,
15 amido or keto groups; and such that the fragment of the ligand defined by the structure $\text{PO-R}^2\text{-OP}$ is chiral. By this definition Applicants intend that the chirality of the diphosphinite ligand arises from the chirality of the parent carbohydrate diol $\text{HO-R}^2\text{-OH}$.

20 Specifically, the process is carried out by employing chiral, nonracemic, O-substituted carbohydrate phosphorus ligands; including particularly pyranose, furanose, disaccharide and oligosaccharide organo-phosphorus ligands. Examples are represented by the
25 formulas IV, V, VI and VII,

10



wherein:

$n = 0-2$;

$m = 0-3$;

R^4 groups are independently H, hydroxy, C_1 to C_{20} hydrocarbyl, alkoxy, aryloxy, O-substituted pyranose or O-substituted furanose;

R^5 groups are independently H, hydroxymethyl (CH_2OH), alkoxymethyl, aryloxy, or $CH_2OP(X)_2$ where X is aryl, alkoxy, or aryloxy;


R^6 groups are independently H, C_1 to C_{20} hydrocarbyl, acyl, or $P(X)_2$ where X is aryl, alkoxy, or aryloxy;

R^7 is H or CH_3 ;


and the sum total of $P(X)_2$ groups present in the O-substituted pyranose, furanose, disaccharide or oligosaccharide organophosphorus ligand is equal to 2.

Applicants also specifically include within the carbohydrate ligand compositions of the invention those carbohydrates containing protective groups. By the term "protective group", Applicants include groups such as ethers and esters which function to provide chiral recognition of the sugar molecule, and further are

commonly employed to protect the sugar molecule from nonselective reactions. Applicants further intend to particularly include disaccharides formed by joining two of the structures shown in formulas IV-VII through an oxygen atom at the anomeric position of the furanose or pyranose ring. Two examples of such dissacharides are shown below, wherein Ph is phenyl and Ac is acetyl.



Most preferably, the chiral, nonracemic, organo-phosphorus ligand is a chiral, nonracemic, O-substituted glucopyranose organophosphorus ligand of the formula VIII,



wherein:

R^8 is H, hydroxy, C_1 to C_{20} hydrocarbyl, alkoxy, or aryloxy;

R⁹ is independently selected from H, C₁ to C₂₀ hydrocarbyl, acyl or P(X)₂, where X is aryl, alkoxy, aryloxy;

5 and the sum total of P(X)₂ groups present in the O-substituted glucopyranose organophosphorus ligand is equal to 2.

Chiral, nonracemic O-substituted carbohydrate derived diolphosphorus ligands can be prepared according to techniques well-known in the art. [J. Organomet. Chem., 159, C29 (1978); Tetrahedron Lett., 1635 (1978); J. Org. Chem., 45, 62 (1980); Bull. Chem. Soc. Jpn., 59, 175 (1986); J. Mol. Catal., 37, 213 (1986); J. Prakt. Chem., 329 (4), 717 (1987)]. In general, diol derivatives containing unprotected hydroxyl groups are treated with a P(R)₂Cl (wherein R may generally be an alkyl, aryl, alkoxy, or aryloxy) reagent, in the presence of a base, such as pyridine or triethylamine, to produce the desired phosphinite or phosphite. Some P(R)₂Cl reagents are commercially available, such as PPh₂Cl (Ph=phenyl). Other P(R)₂Cl reagents, where R = aryl or alkyl, can be prepared by two methods. Method A involves the reaction of (amino)dichlorophosphines such as Et₂NPCl₂ with RMgBr followed by reaction with HCl [J. Am. Chem. Soc. 2148, 82, (1960)]; J. Am. Chem. Soc. 80, 1107 (1958). Alternatively, treatment of readily available dialkyl phosphites, such as dibutyl phosphite, HP(O)(OBu)₂, with RMgBr followed by reaction with PCl₃ provides P(R)₂Cl derivatives. [J. Am. Chem. Soc. 73, 4101 (1951); J. Am. Chem. Soc., 74, 5418 (1952)]; J. Org. Chem., 31, 1206 (1966). P(R)₂Cl reagents, where R=alkoxy or aryloxy, can be prepared in two steps by treatment of P(NEt₂)₃ with ROH to generate P(OR)₂(NEt₂), followed by treatment with CH₃COCl to generate P(OR)₂Cl. Illustrative preparations are provided below.

For all embodiments of the invention the chiral, nonracemic metal hydroformylation catalyst may be prepared by mixing the metal source and the chiral, nonracemic, organophosphorus ligand, preferably in a suitable organic solvent under an inert atmosphere such as N₂ or Ar in a temperature range from 0°C to 120°C, preferably in a temperature range from 0°C to 80°C. The metal compound may be used in this solution or the metal compound can be obtained in the pure form upon removal of the solvent.

The molar ratio of chiral, nonracemic, organophosphorus ligand to the metal may vary between 1:1 to 10:1, preferably between 1:1 to 1.2:1.

The molar ratio of metal complex to vinyl compound may vary between 0.0001:1 to 1:1, preferably between 0.0025:1 to 0.05:1.

The vinyl compound starting material, which is represented by the formula R-CH=CH₂ may be dissolved in any organic solvent compatible with the reagents employed, preferably a nonpolar solvent (nonpolar solvents generally provide higher ee's) such as, but not limited to, benzene, hexane, or triethylsilane. In the case of a liquid substrate, the substrate itself can serve as the solvent.

The H₂ and CO can be provided by contacting the reaction mixture with the gases and can be provided in mixtures of H₂ to CO ranging from 10:1 to 1:10.


A preferred 1:1 mixture of H₂ and CO can be conveniently prepared or purchased commercially. The pressures of H₂ and CO under which the hydroformylation reactions can be conducted range generally from 100-2400 psi (1 psi= 6.9 kPa), with 500-1000 psi being the preferred pressure. The pressure employed is not critical; for example, in the case of the hydroformylation of 2-vinylnaphthalene in hexane at room

temperature using catalyst 1, the following ee's were obtained with the pressure of H₂ and CO (1:1) indicated: 49% (500 psi), 51% (1600 psi), and 31% (2400 psi).

5 The hydroformylation reaction is preferably carried out over a temperature range from 20° to 80°C, most preferably 25° to 30°C. Applicants note that the observed overall selectivity (ee's and branched to linear ratios) generally decreases as the temperature is increased. For example, the hydroformylation of
10 2-vinylnaphthalene in benzene at 1600 psi provides an ee of 10% when the reaction is performed at room temperature and 1% at 70°C. The branched to linear ratio in this case falls from 21:1 (room temperature) to 12:1 (70°C). In addition, the ee of the hydro-
15 formylation of 2-vinylnaphthalene decreases from 12% to 11% when the reaction is conducted in THF at room temperature versus 70°C. The branched to linear ratio also decreases in the latter example from 40:1 (room temperature) to 10:1 (70°C).

20 The enantioselective hydroformylation reactions are generally complete within 18-48 hours. In the case of platinum catalysts, the presence of a Lewis acid such as tin (II) chloride is typically preferred.


To demonstrate a preferred mode of the invention
25 which produces a particularly useful product, preparation of optically active (S)-(-)-2-(6-methoxy-2-naphthalene)propanal, can be achieved. The catalyst composition comprises a cationic rhodium (I) compound and the ligand of formula II wherein each R¹ is the aryl
30 group 3,5-bis(trifluoromethyl)phenyl and R² is the O-substituted β-D-glucopyranose derivative of the formula IX, the starting vinyl compound is 6-methoxy-2-vinylnaphthalene, and the preferred source of the rhodium(I) species is (COD)₂RhBF₄.



For the preparation of optically active (S)-(-)-2-(6-methoxy-2-naphthalene)propanal, the enantioselective hydroformylation is preferably carried out at 25°C under 500 p.s.i. pressure of a 1:1 mixture of H₂ and CO. A mixture of 6-methoxy-2-vinylnaphthalene and the chiral rhodium complex in a non-polar organic solvent such as benzene, hexane, or triethylsilane is shaken at room temperature for 18 h. In this preferred embodiment a molar ratio between 0.0025:1 to 0.05:1 of rhodium catalyst to vinyl compound is preferred. A molar ratio between 1:1 to 1:1.2 of metal to organophosphorus ligand is preferred.

Using these preferred conditions an ee between 35-72% of the S enantiomer of 2-(2-naphthyl)propanal and a yield between 75-95% will generally be obtained. Isolation of the product aldehyde can be achieved by flash column chromatography of the reaction mixture on silica gel using 10% diethyl ether/hexane as eluent.

20 General Procedures for the Preparation of Chiral, O-Substituted Carbohydrate Phosphinite and Phosphite Ligands.



Ligand 1; $R^1 = 3,5-(CF_3)_2C_6H_3$

Ligand 2; $R^1 = Ph$ (phenyl)


Ligand 3; $R^1 = 4-FC_6H_4$

Ligand 4; $R^1 = (4-CF_3)C_6H_4$

5 The ligands 1 through 4 above were prepared according to the method previously reported in U.S. Patent 5,175,335, which is hereby generally incorporated by reference. However, a modified procedure for the synthesis of Ar_2PCl (ligand 5) was developed (see
10 below). All reactions were carried out under a N_2 atmosphere using standard Schlenk techniques or a Vacuum Atmospheres Co. Drybox. Solvents were distilled and degassed prior to use.

15 Example of Modified Procedure for Synthesis of
 Ligand 5, Ar_2PCl species; Di-[(3,5-bis-trifluoro-
 methyl)phenyl]chlorophosphine

 A 1.0 M solution of (3,5-bis-trifluoromethyl)-phenylmagnesium bromide was prepared in the dry box by
20 slow addition of 18.5 g (60 mmol) of (3,5-bis-trifluoromethyl)bromobenzene in 40 mL of THF to a slurry of Mg turnings in 20 mL of THF. After 1 h, this solution was added slowly to a solution of 5.0 g (29 mmol) of Et_2NPCl_2 in 30 mL of THF at $0^\circ C$. After 2 h, the mixture
25 was concentrated in vacuo. Cyclohexane (100 mL) was added and the mixture was filtered to provide a solution of [di-3,5-bis(trifluoromethyl)-phenyl](diethylamino)-phosphine. Dry HCl was passed through this solution for 1 h. After filtration and concentration, 12.4 g (88%)
30 of di-[(3,5-bis-trifluoromethyl)-phenyl]chlorophosphine was collected as a white solid. ^{31}P (C_6D_6) δ 70.4, s. 1H (C_6D_6) δ 7.54 (s, 1) 7.66 (d, 1, $J = 7$ Hz).

Synthesis of Catalysts

Catalyst 1: Ar = 3,5-(CF₃)₂C₆H₃; A = BF₄

(Phenyl 2,3-bis-O-[[di-(3,5-bis-trifluoro-methyl)-phenyl]phosphino]-4,6-benzylidene-β-D-glucopyranoside)-(1,5-cyclooctadienyl) rhodium(I) tetrafluoroborate.

To a solution of 89 mg (0.22 mmol) of (COD)₂RhBF₄ in 2 mL of CH₂Cl₂ was added a solution of 300 mg (0.24 mmol) of ligand 1 in 2 mL of CH₂Cl₂. After 30 min, the mixture was concentrated to give a 2:1 mixture of complexes of catalyst 1. Major complex showed: ³¹P NMR δ 126.7 (dd, 1, J = 40, 184 Hz), 123.0 (dd, 1, J = 40, 179 Hz); Minor complex showed: ³¹P NMR δ 139.7 (dd, 1, J = 48, 234 Hz), 124.1 (dd, 1, J = 48, 240 Hz).

Catalyst 2: Ar = Ph; A = BF₄

(Phenyl 2,3-bis-O-diphenylphosphino)-4,6-benzylidene-β-D-glucopyranoside)-(1,5-cyclooctadienyl) rhodium(I) tetrafluoroborate.

In a similar fashion, catalyst 2 was prepared from 31 mg of (COD)₂RhBF₄ and 57 mg of ligand 1. Catalyst 2 showed: ³¹P NMR δ 134.5 (dd, 1, J = 28, 179 Hz), 137.5 (dd, 1, J = 28, 178 Hz).

Catalyst 3: Ar = 3,5-(CF₃)₂C₆H₃; A = OSO₂CF₃
(Phenyl 2,3-bis-O-([di-(3,5-bis-trifluoro-methyl)-
phenyl]phosphino)-4,6-benzylidene-β-D-glucopyranoside)-
1,5-cyclooctadienyl)rhodium (I) trifluoromethane-
5 sulfonate.

In a similar fashion, catalyst 3 was prepared from
40 mg of (COD)₂Rh(OSO₂CF₃) and 100 mg of ligand 1.
Catalyst 3 showed: ³¹P NMR δ 122.6 (dd, 1, J = 44, 176
Hz), 126.0 (dd, 1, J = 44, 166 Hz).

Catalyst 4: Ar = 4-FC₆H₄; A = BF₄
(Phenyl {2,3-bis-O-[di-(4-fluorophenyl)-phosphino]}-4,6-
benzylidene-β-D-glucopyranoside) (1,5-cyclooctadienyl)-
rhodium (I) tetrafluoroborate.

In a similar fashion, catalyst 4 was prepared from
49 mg (0.121 mmol) of (COD)₂RhBF₄ and 100 mg
(0.128 mmol) of 3. Catalyst 4 showed: ³¹P (CDCl₃) δ
131.3 (dd, 1, J = 30, 181), 136.0 (dd, 1, J = 32, 179
Hz).


Catalyst 5: Ar = 3,5-(CF₃)₂C₆H₃; A = SbF₆
(Phenyl (2,3-bis-O-([di-(3,5-bis-trifluoro-methyl)-
phenyl]phosphino)-4,6-benzylidene-β-D-glucopyranoside)-
(1,5-cyclooctadienyl) rhodium (I) hexafluoroantimonate.

In a similar fashion, catalyst 5 was prepared from
42 mg (0.076 mmol) of (COD)₂RhSbF₆ and 100 mg
(0.080 mmol) of ligand 1. A ca. 4:1 mixture of
complexes of catalyst 5 was obtained. Major complex
showed: ³¹P NMR δ 126.6 (dd, 1 J = 34, 182 Hz), 130.7
(dd, 1, J = 34, 186 Hz); Minor complex showed: ³¹P NMR δ
125.6 (dd, 1, J = 44, 239 Hz), 139.8 (dd, 1, J = 44, 236
Hz).

Catalyst 6: Ar = (4-CF₃)C₆H₄; A = BF₄

(Phenyl (2,3-bis-O-([di-(4-trifluoromethyl)-phenyl]-phosphino)-4,6-benzylidene-β-D-glucopyranoside)-(1,5-cyclooctadienyl) rhodium (I) tetrafluoroborate.

- 5 A solution of catalyst 6 was prepared in situ from ligand 4 and (COD)₂RhBF₄.



Catalyst 7: Ar = Ph

- 10 {(Phenyl [2,3-bis-O-(diphenylphosphino)]-4,6-benzylidene-β-D-glucopyranoside)platinum (II) dichloride.

To a solution of 50 mg (0.11 mmol) of (PhCN)₂PtCl₂ in 3 mL of benzene was added a solution of 90 mg (0.13 mmol) of ligand 2 in 2 mL of benzene. After 1 h, the mixture was filtered through a glass wool plug and concentrated to give catalyst 7: ³¹P NMR δ 95.1 (d, 1, J = 11 Hz), 94.4 (d, 1, J = 11 Hz).

Catalyst 8: Ar = 3,5-(CF₃)₂C₆H₃

- 20 {(Phenyl [di-(3,5-bis-trifluoromethyl)-phenyl]-phosphino)-4,6-benzylidene-β-D-glucopyranoside}platinum (II) dichloride.

A solution of catalyst 8 was prepared in situ from ligand 1 and (PhCN)₂PtCl₂.

- 25 Catalyst 9: Chiral Iridium Catalyst

Preparation of Chiral Iridium Catalyst: To a solution of 50 mg (0.074 mmol) of [Ir(COD)Cl]₂ in 5 mL of CH₃CN was added a solution of 198 mg (0.158 mmol) of ligand

1 in 5 mL of CH₃CN. After 1 h, the solution was concentrated to give catalyst 9 as a yellow-orange powder.

5 Catalyst 10: Chiral Cobalt Catalyst

Chiral cobalt catalyst 10 was generated in situ from 4 mg of Co₂(CO)₈ and 22 mg of ligand 1 in 2 mL of THF.

EXAMPLES 1-36

General Methods for Enantioselective

10 Hydroformylation of Vinyl Compounds, for Examples 1-36

Results and reaction conditions for the enantioselective hydroformylation of vinyl compounds are shown in Tables I-IV. With a few exceptions which are noted, the hydroformylations were conducted by shaking a solution of the vinyl compound under a pressure of H₂ and CO at the noted temperature. Conversions were determined by integration of the formyl protons of the product aldehydes and the vinyl protons of the vinyl compounds in the ¹H NMR spectrum of the crude product mixture. EE's were determined after reduction of the crude mixture with lithium aluminum hydride to generate the corresponding primary alcohols by HPLC using a Bakerbond Chiral DNBPG Chiralcel OJ or OB column (J. T. Baker, Phillipsburg, NJ): 2-5% i-PrOH/Hexane, 1 mL/min., 40°C. HPLC samples were passed through a short pad of silica gel and eluted with a hexane/Et₂O gradient prior to analysis. In some instances ee's were determined by ¹H NMR using chiral shift reagent Eu(hfc)₃ or by gas chromatographic separation using a Chiraldex G-TA capillary column (Aztec, Whippany, NJ). EE values refer to an excess of the S-enantiomer. Enriched samples of the S-alcohols from the hydroformylation of 2-vinylnapthalene and 6-methoxy-2-vinylnapthalene were obtained by diisobutylaluminum hydride reduction of the corresponding nitrile [J. Am. Chem. Soc. 114, 6265

(1992)] followed by lithium aluminum hydride reduction. The relative assignments of the other hydroformylation products were assigned by analogy to be enriched with the S-enantiomer.

5

Typical Procedure For Hydroformylation Reactions:

In the dry box a 140 mL glass liner was charged with 2.0 mmol of vinyl compound, 0.003 mmol of catalyst/cocatalyst, and 10 mL of solvent. This liner was placed in a shaker tube and pressurized with a 1:1 mixture of hydrogen and carbon monoxide. After venting, the mixture was pressurized with the hydrogen, carbon monoxide mixture and shaken at the given temperature. Upon completion, the mixture was vented and the glass liner was removed from the shaker tube. The solution was filtered through celite and concentrated. Percent conversions were determined by ^1H NMR integration of the resonances corresponding to the starting vinyl compound and product aldehydes. The branched aldehyde to linear aldehyde ratio (b/l) was determined by ^1H NMR integration of the resonances corresponding to the aldehyde protons or by GC.

The crude mixture was dissolved in 10 mL of ether and 50 mg (1.3 mmol) of LiAlH_4 was added. After 3 h, water (4 drops), 15% NaOH (4 drops) and water (12 drops) was added carefully. After 30 min. Na_2SO_4 was added and the mixture was filtered and concentrated. Flash chromatography on silica gel using 40% ether/hexane as eluant gave 2-substituted-1-propanols. The ees of these alcohols were determined by analysis on either an OB or OJ chiralcel HPLC column using an *i*-propanol/ hexane mixture as eluant.

EXAMPLES 1-12TABLES I AND II

Hydroformylation of 2-Vinylnapthalene using
Catalyst Composition 1 or 2

5

EXAMPLE 1

Following the general procedure, a mixture of
300 mg (2.0 mmol) of 2-vinylnapthalene and 5 mg of
catalyst 1 in 10 mL of Et₃SiH was pressurized at room
temperature to 1600 psi using the H₂/CO mixture. After
10 18 h, the mixture was manipulated as described:
conversion = 20%; b/l = 22; ee = 72%.

EXAMPLE 2

The reaction was carried out similar to Example 1,
using hexane as solvent and 500 psi of H₂/CO provided
15 the following: conversion = 100%; b/l = 20; ee = 50%.

EXAMPLE 3

The reaction was carried out similar to Example 1,
using hexane and triethylformate as solvent and 1600 psi
of H₂/CO provided the following: conversion = 85%; b/l
20 = 20; ee = 17%. In this case the ee was determined on
the diethyl acetal resulting from in situ trapping of
2-(2-naphthyl)-1-propanal.

Examples 4-9 were performed in a similar fashion
under the conditions described in Table I.

25 Examples 10-12 were performed in a similar manner
using catalyst 2 under the conditions described in
Table II.

TABLE I

Asymmetric Hydroformylation
of 2-Vinylnaphthalene Using Catalyst 1

<u>Ex.</u>	<u>Solvent</u>	<u>Pressure (psi)</u>	<u>Temp. (°C)</u>	<u>b/l</u>	<u>Conv. (%)</u>	<u>ee (%)</u>
1	Et ₃ SiH	1600	ambient	22	20	72
2	Hexane	500	ambient	20	100	49
3	Hexane + CH(OEt) ₃	1600	ambient	20	85	17
4	Hexane	1600	ambient	26	53	51
5	Hexane	2400	ambient	24	80	31
6	Benzene	1600	ambient	33	43	38
7	THF	1600	ambient	40	71	12
8	THF	1600	70	11	100	10
9	THF	2400	ambient	-	100	7

TABLE II

Asymmetric Hydroformylation
of 2-Vinylnaphthalene Using Catalyst 2

<u>Ex.</u>	<u>Solvent</u>	<u>Pressure (psi)</u>	<u>Temp. (°C)</u>	<u>b/l</u>	<u>Conv. (%)</u>	<u>ee (%)</u>
10	Benzene	1600	ambient	21	18	10
11	Benzene	1600	70	12	100	1
12	THF	1600	70	10	100	11

EXAMPLES 13-26TABLE III

Examples 13-26 were performed in a similar fashion using the appropriate vinyl compound and catalyst as indicated. Abbreviations are as follows 6-methoxy-2-vinylnaphthalene (MVN), styrene (ST), 4-isobutylstyrene (IBS), 4-methylstyrene (4-MeST), vinyl acetate (VA).

L' =




TABLE III

Other Asymmetric Hydroformylation
Reactions Using L'¹Rh(COD)A Complexes

Ex.	Substrate	R ¹	A	Solvent	b/l	Conv. (%)	ee (%)
13	VN	(4-CF ₃)C ₆ H ₄	BF ₄	Hexane	10	40	10
14	MVN	(4-CF ₃)C ₆ H ₄	BF ₄	Hexane	8	10	15
15	MVN	4-FC ₆ H ₄	BF ₄	Hexane	7	10	5
16	VN	3,5-(CF ₃) ₂ C ₆ H ₃	SbF ₆	Hexane	18	100	19
17	ST	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Hexane	29	50	24
18	ST	3,5-(CF ₃) ₂ C ₆ H ₃	SbF ₆	Hexane	19	63	17
19	ST	3,5-(CF ₃) ₂ C ₆ H ₃	OTf	Hexane	21	53	14
20	4-MeST	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Hexane	16	43	30
21	IBS	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Et ₃ SiH	16	1	43
22	MVN	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Hexane	15	60	44
23	MVN	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Et ₃ SiH	20	5	70
24	MVN	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	i-Pr ₃ SiH	14	14	16
25	VA	3,5-(CF ₃) ₂ C ₆ H ₃	BF ₄	Hexane	12	10	14
26	VA	3,5-(CF ₃) ₂ C ₆ H ₃	OTf	Hexane	9	10	11

EXAMPLES 27-30

Asymmetric Hydroformylation of
6-Methoxy-2-vinylnaphthalene and
2-Vinylnaphthalene Using Chiral Platinum Catalysts

5

EXAMPLE 27

Following the general procedure, a mixture of
400 mg (2.2 mmol) of 6-methoxy-2-vinylnaphthalene, 3 mg
(0.016 mmol) of SnCl₂, 2 mg (0.016 mmol) of 4-methoxy-
phenol and 10 mg (0.001 mmol) of {(phenyl [2,3-bis-O-
(diphenylphosphino)]-4,6-benzylidene-β-D-glucopyranoside}platinum (II) dichloride (catalyst 7) in
10 mL of benzene was pressurized at ambient temperature
to 2400 psi using the H₂/CO mixture. After 44 h, the
mixture was manipulated as described: conversion =
100%; b/l = 0.7; ee = 9%.

EXAMPLE 28

The reaction was carried out similar to Example 31, except performing the reaction at 60°C which provided the following: conversion = 100%; b/l = 1.3; ee = 9%.

5

EXAMPLE 29

The reaction was carried out similar to Example 31, using benzene and triethylformate as solvent, which provided the following: % conversion = 80%; b/l = 1.5; ee = 9%. In this case the ee was determined on the diethyl acetal resulting from in situ trapping of 2-(2-naphthyl)-1-propanal.

10

EXAMPLE 30

In this case a benzene solution of chiral platinum catalyst, (phenyl 2,3-bis-O-([di-(3,5-bis-trifluoro-methyl)-phenyl]phosphino)-4,6-benzylidene-β-D-glucopyranoside)platinum(II) dichloride (catalyst 8), was generated from 24 mg (0.02 mmol) of 1 and 7 mg (0.02 mmol) of dichlorobis(benzonitrile)platinum (II). This catalyst solution, 300 mg (2.0 mmol) of 2-vinyl-naphthalene, 6 mg (0.032 mmol) of SnCl₂, 2 mg (0.016 mmol) of 4-methoxyphenol, and 8 mL of benzene was placed in a glass liner. The reaction was performed at 60°C and 2400 psi pressure of H₂/CO for 18 h to provide the following: conversion = 90%; b/l = 2.7; ee = 5%.

20

EXAMPLE 31

Asymmetric Hydroformylation of
6-Methoxy-2-vinylnaphthalene Using Chiral Iridium
Catalysts

The reaction was performed in a similar manner to the rhodium-catalyzed hydroformylations; 300 mg of MVN, 5 mg of chiral iridium catalyst 9, and 10 mL of THF. This experiment was conducted at 80°C using 3000 psi of H₂/CO to provide the following: conversion 37%; b/l = 10; ee = 30%.

30

EXAMPLE 32

Asymmetric Hydroformylation of
6-Methoxy-2-vinylnapthalene Using Chiral Cobalt
Catalysts

- 5 This reaction was performed in a similar manner to the rhodium-catalyzed hydroformylations, except the chiral cobalt catalyst 10 was generated in situ from 4 mg of $\text{Co}_2(\text{CO})_8$ and 22 mg of 5 in 2 mL of THF. The reaction was performed using this catalyst solution,
- 10 300 mg of MVN, and 8 mL of THF. The reaction was conducted at 120°C and 5000 psi of H_2/CO to provide the following: conversion = 10%; b/l = 3; ee = 1%.

WHAT IS CLAIMED IS:

1. A process for enantioselective hydroformylation comprising: reacting a vinyl compound of formula I




with a source of CO and H₂, in the presence of a catalyst composition comprising one or more metals selected from the group consisting of Co, Rh, Ir, and Pt, and a chiral, nonracemic ligand of formula II

(R¹)₂-P-O-R²-O-P-(R¹)₂

II

to produce a nonracemic hydroformylated product of formula III



15

wherein

R is a C₁ to C₂₀ carboalkoxy, a C₁ to C₄₀ aromatic hydrocarbyl, or a C₁ to C₄₀ heterocyclic radical; each optionally substituted with one or more halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups;

each R¹ is an electron-withdrawing group comprising an aromatic hydrocarbyl substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkylsulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups;

20

25

or a heteroaromatic optionally substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups;

R² is a C₄ to C₄₀ dideoxy carbohydrate optionally substituted with one or more hydrocarbyl, halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups.


2. The process of Claim 1, wherein the metal is Rh.

3. The process of Claim 1, wherein the reaction is carried out at a pressure of about 100 psi to about 2400 psi under inert atmosphere and at a temperature of about 20°C to about 80°C.

4. The process of Claim 1, wherein the molar ratio of ligand to metal is about 1 to 1, to about 1.2 to 1, and wherein the molar ratio of metal to starting vinyl compound is about 0.0025 to 1, to about 0.05 to 1.

5. The process of Claim 1 wherein the H₂ and CO are provided in a ratio of about 10 to 1 to about 1 to 10.

6. The process of Claim 1, wherein the starting vinyl compound is 6-methoxy-2-vinyl-naphthalene, the metal is Rh, each R¹ is 3,5-bis(trifluoromethyl)phenyl, R² is the compound of formula IX, and the hydroformylated product is (S)-(-)-2-(6-methoxy-2-naphthalene)propanal.



IX

7. The process of Claim 1, wherein the metal is Pt and further comprising wherein the reaction is carried out in the presence of a Lewis acid.

8. The process of Claim 1 wherein R² is
5 4,6-benzylidene-β-D-glucopyranoside.

9. The process of Claim 1 wherein the ligand of formula II is phenyl 2,3-bis-O-([di-(3,5-bis-trifluoromethyl)-phenyl]phosphino)-4,6-benzylidene-β-D-glucopyranoside.

10 10. A catalyst composition comprising one or more metals selected from the group consisting of Co, Rh, Ir and Pt, and a chiral nonracemic ligand of formula II



II

wherein

15 each R¹ is an electron-withdrawing group comprising an aromatic hydrocarbyl substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkylsulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups; or
20 a heteroaromatic optionally substituted with one or more halo, halogen-substituted alkyl groups, cyano, alkyl sulfonyl, carboalkoxy, quaternary ammonium, nitro, amido or keto groups;
25 and

R² is a C₄ to C₄₀ dideoxy carbohydrate; optionally substituted with one or more hydrocarbyl, halo, alkoxy, carboalkoxy, hydroxy, amido or keto groups.

30 11. The composition of Claim 10, wherein the metal is Rh.

12. The composition of Claim 10, wherein the ratio of ligand to metal is from about 1 to 1, to about 1.2 to 1.

13. The composition of Claim 10, wherein R¹ is
5 3,5-(CF₃)₂C₆H₃.

14. The composition of Claim 10, wherein R² is 4,6-benzylidene-β-D-glycopyranoside.

15. The composition of Claim 10, wherein the metal is Rh and the ligand of formula II is phenyl 2,3-bis-O-
10 {[di-(3,5-bis-trifluoromethyl)-phenyl]phosphino}-4,6-benzylidene-β-D-glucopyranoside.

INTERNATIONAL SEARCH REPORT

Intern: 1 Application No
PCT/US 94/09461

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C45/50 B01J31/18 C07C47/277 C07C47/23 C07C45/49

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07C B01J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DD,A,280 473 (AKADEMIE DER WISSENSCHAFT DER DDR) 11 July 1990 cited in the application see page 2, paragraph 2; claims	1,10
X	TETRAHEDRON: ASYMMETRY, vol.4, no.3, March 1993, OXFORD GB pages 369 - 382 R SELKE ET AL see page 371	10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

28 October 1994

Date of mailing of the international search report

- 9. 11. 94

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Heywood, C

INTERNATIONAL SEARCH REPORT

Internat'l Application No

PCT/US 94/09461

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 109, no. 25, 19 December 1988, Columbus, Ohio, US; abstract no. 231476v, page 911 ; see abstract & J.PRAKT.CHEM., vol.329, no.4, 1987 pages 717 - 724 ----	10
A	JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol.114, 1992, WASHINGTON, DC US pages 6265 - 6266 T RAJANBABU ET AL see the whole document ----	1,10
A	FR,A,2 208 872 (INSTITUT FRANCAIS DU PETROLE,DES CARBURANTS ET LUBRIFIANTS) 28 June 1974 see the whole document ----	1
P,X	TETRAHEDRON LETTERS, vol.35, no.25, June 1994, OXFORD GB pages 4295 - 4298 T RAJANBABU ET AL see the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/09461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DD-A-280473		NONE	
FR-A-2208872	28-06-74	NONE	